



## Environmental Monitoring and Reporting Canning River Estuary ~ Weekly profile

by  
Aquatic Science Branch, Department of Water and Environmental Regulation

Date: Wednesday 30<sup>th</sup> January 2019

Weather & tide conditions: Conditions were fine with 0% cloud cover and a predominantly south-easterly breeze of up to 4 knots. The tide was flowing during the sampling period. Gosnells recorded 4.6 mm of rainfall during the week prior to the morning of sampling (Bureau of Meteorology).

Oxygenation: The Nicholson Road and Bacon Street oxygenation plants were operational the week prior to sampling.

Canning Estuary (downstream of Kent St Weir): The Canning Estuary was saline at SCB2 and SAL, saline to brackish at RIV and brackish at CASMID. Water was well oxygenated to oxygenated throughout, with the exception of low oxygen bottom water at RIV and CASMID. Water temperature ranged from 24.2 to 30.2 ° C.

Lower Canning River (upstream of Kent St Weir): The Lower Canning River was fresh throughout all sites. Water was mostly well oxygenated to oxygenated throughout all sites, with the exception of low oxygen surface water at ELL, near anoxic bottom water at ELL, KEN, BAC and PAC, and low oxygen bottom water at KS7. Water temperature ranged from 21.5 to 26.3 ° C.

**Definitions:**

Salinity – fresh <5ppt, brackish 5-25ppt, saline 25-35ppt, hypersaline >35ppt

Dissolved oxygen – well oxygenated >6 mg/L  
oxygenated >4 to 6 mg/L, low oxygen >2 to 4 mg/L,  
hypoxic >0.5 to 2.0 mg/L, near anoxic <0.5 mg/L

# Canning River Estuary - Physical-Chemical Profile - 30th January 2019

Prepared by Aquatic Science Branch, Department of Water and Environmental Regulation

